



**Elemental Enzymes Ag & Turf, LLC**  
 Vismax® Row Crop  
 EPA Reg. No. 92188-4

**FIFRA Section 2(ee) Recommendation**  
 Issue Date: [March 1, 2026](#)  
 Expiration Date: [March 1, 2029](#)

FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY

**Directions For Use**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which are recommended by Elemental Enzymes Ag & Turf, LLC. These directions may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product label.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

This recommendation must be in the possession of the user at the time of pesticide application.

Crop	Targeted Diseases	Application Method	Application Rate	Application Instructions
Wheat	<b>Fusarium Head Blight</b> ( <i>Fusarium graminearum</i> )	Foliar	2.0 - 4.0 fl oz/acre	Apply up to 2 times per cropping season using ground or aerial equipment at vegetative and reproductive stages as a preventative treatment, or at first sign of disease.
Corn	<b>Gray Leaf Spot</b> ( <i>Cercospora zeae-maydis</i> ) <b>Northern Leaf Blight</b> ( <i>Exserohilum turcicum</i> ) <b>Rusts</b> ( <i>Puccinia</i> spp.) <b>Tar Spot</b> ( <i>Phyllachora maydis</i> )	Foliar	2.0 - 4.0 fl oz/acre	Apply up to 2 times per cropping season using ground or aerial equipment at vegetative and reproductive stages starting at V4 (4 leaf collars) as a preventative treatment, or at first sign of disease.
Soybean	<b>Frogeye Leaf Spot</b> ( <i>Cercospora sojina</i> ) <b>Septoria Brown Spot</b> ( <i>Septoria glycines</i> )	Foliar	2.0 - 4.0 fl oz/acre	Apply up to 2 times per cropping season using ground or aerial equipment at vegetative and reproductive stages as a preventative treatment, or at first sign of disease.

